

# Disparities in Cancer Screening in Individuals With a Family History of Breast or Colorectal Cancer

Ninez A. Ponce, MPP, PhD<sup>1,2</sup>; Jennifer Tsui, MPH<sup>1,2</sup>; Sara J. Knight, PhD<sup>3,4,5,6</sup>; Aimee Afable-Munsuz, PhD<sup>6,7</sup>; Uri Ladabaum, MD, MS<sup>8</sup>; Robert A. Hiatt, MD, PhD<sup>6,9</sup>; and Jennifer S. Haas, MD, MSc<sup>10</sup>

**BACKGROUND:** Understanding racial/ethnic disparities in cancer screening by family history risk could identify critical opportunities for patient and provider interventions tailored to specific racial/ethnic groups. The authors evaluated whether breast cancer (BC) and colorectal cancer (CRC) disparities varied by family history risk using a large, multiethnic population-based survey. **METHODS:** By using the 2005 California Health Interview Survey, BC and CRC screening were evaluated separately with weighted multivariate regression analyses, and stratified by family history risk. Screening was defined for BC as mammogram within the past 2 years for women aged 40 to 64 years; for CRC, screening was defined as annual fecal occult blood test, sigmoidoscopy within the past 5 years, or colonoscopy within the past 10 years for adults aged 50 to 64 years. **RESULTS:** The authors found no significant BC screening disparities by race/ethnicity or income in the family history risk groups. Racial/ethnic disparities were more evident in CRC screening, and the Latino-white gap widened among individuals with family history risk. Among adults with a family history for CRC, the magnitude of the Latino-white difference in CRC screening (odds ratio [OR], 0.28; 95% confidence interval [CI], 0.11-0.60) was more substantial than that for individuals with no family history (OR, 0.74; 95% CI, 0.59-0.92). **CONCLUSIONS:** Knowledge of their family history widened the Latino-white gap in CRC screening among adults. More aggressive interventions that enhance the communication between Latinos and their physicians about family history and cancer risk could reduce the substantial Latino-white screening disparity in Latinos most susceptible to CRC. *Cancer* 2011;000:000-000. © 2011 American Cancer Society.

**KEYWORDS:** family history, cancer screening, cancer disparities, breast cancer, colorectal cancer.

## INTRODUCTION

**Having** a family history of breast cancer (BC) or colorectal cancer (CRC) is associated with an increased risk of developing these cancers compared with those with no family history.<sup>1-3</sup> BC and CRC are among the most common types of cancers as well as among the leading causes of cancer death in the United States, and both have evidence-based recommendations for screening and prevention.<sup>4</sup> Nationally, among the estimated 8% of the population who report a family history of BC and 5% who report a family history of CRC, screening rates for these cancers are higher than among average-risk individuals.<sup>5</sup> Although on average, knowledge of family history clearly motivates individuals to receive cancer screening, less is known regarding whether there are racial/ethnic disparities in screening among this higher-risk group, as indicated in many studies on average-risk populations.<sup>6-10</sup> Understanding racial/ethnic disparities in cancer screening by family history risk could identify critical opportunities for patient and provider interventions tailored to specific racial/ethnic groups. Focusing on individuals who are most susceptible to breast and colorectal cancers could effectively target resources to reduce the disproportionate burden of cancer incidence and mortality in racial/ethnic minority populations.<sup>6,11,12</sup>

**Corresponding author:** Ninéz A. Ponce, MPP, PhD, Department of Health Services and Division of Cancer Prevention & Control Research, UCLA School of Public Health & Jonsson Comprehensive Cancer Center, 650 Charles Young Drive South, Los Angeles, CA 90095-1772; Fax: (310) 825-3317; nponce@ucla.edu

<sup>1</sup>Department of Health Services, University of California at Los Angeles, Los Angeles, California; <sup>2</sup>Jonsson Comprehensive Cancer Center, University of California at Los Angeles, Los Angeles, California; <sup>3</sup>San Francisco Veterans Affairs Medical Center, San Francisco, California; <sup>4</sup>Department of Psychiatry, University of California at San Francisco, San Francisco, California; <sup>5</sup>Department of Urology, University of California at San Francisco, San Francisco, California; <sup>6</sup>Helen Diller Family Comprehensive Cancer Center, University of California at San Francisco, San Francisco, California; <sup>7</sup>Division of General Internal Medicine, Department of Medicine, University of California at San Francisco, San Francisco, California; <sup>8</sup>Division of Gastroenterology, Stanford University School of Medicine, Stanford, California; <sup>9</sup>Department of Epidemiology and Biostatistics, University of California at San Francisco, San Francisco, California; <sup>10</sup>Brigham and Women's Hospital/Harvard University, Boston, Massachusetts

**DOI:** 10.1002/cncr.26480, **Received:** January 14, 2011; **Revised:** June 29, 2011; **Accepted:** June 29, 2011, **Published online** in Wiley Online Library (wileyonlinelibrary.com)

Among adults with a family history of cancer, racial/ethnic differences in individual and physician factors, and in the communication between individuals and their physicians, may result in variations in preventive screening behavior.<sup>13</sup> Recent evidence indicates that racial/ethnic minorities are less likely than non-Latino whites to recognize family history as a potential risk factor for familial cancers.<sup>14</sup> This suggests that racial/ethnic minorities are underestimating their cancer risk compared with non-Latino whites, and/or their physicians are not appropriately counseling them on their risk. Even if adults are aware of their family history as a risk factor, risk awareness could manifest differently across racial/ethnic groups; some may vigilantly seek preventive screening, but others may avoid screening because of fear,<sup>15,16</sup> anxiety, and distress.<sup>17</sup> Moreover, researchers have suggested that compared with non-Latino whites, some racial/ethnic minorities are less likely to discuss experiences of cancer because of cultural stigma and taboos surrounding disclosure of cancer experiences.<sup>18,19</sup> There is also evidence that racial/ethnic minorities compared with non-Latino whites are less likely to receive a provider recommendation for cancer screening.<sup>20,21</sup> We suspect that many of these factors, already established among average-risk groups, would be amplified among individuals with a family history, especially with the recent population-based finding of lower perceived cancer risk among racial/ethnic minorities with a family history.<sup>14</sup>

We hypothesized that racial/ethnic screening disparities among individuals reporting family history of breast/ovarian cancers or CRC will be wider than the racial/ethnic screening disparities in screening among individuals reporting no family history of these cancers. By using a population-based survey in California, a diverse state, we examined the nonelderly California adult population to evaluate how family history affects screening behavior. We separately examined racial/ethnic disparities in screening behavior by family history risk.

## MATERIALS AND METHODS

### **Data Source**

We used data from adult respondents who participated in the 2005 California Health Interview Survey (CHIS). CHIS is a population-based random-digit dial telephone survey conducted every other year since 2001 among residents of California.<sup>22</sup> CHIS employs a multistage sampling design to ensure that minority subgroups and rural populations in California are represented in the data.<sup>22</sup>

The 2005 CHIS was administered in 5 languages: English, Spanish, Chinese (Cantonese and Mandarin), Korean, and Vietnamese. Data were collected on a range of health and health-related topics, including cancer screening behavior, personal history of cancer, health status, insurance status, mental health, and health behaviors. The response rate for the CHIS 2005 adult sample was 26.9%, comparable to other large population-based telephone surveys, including the 2005 California Behavioral Risk Factor Surveillance System survey.<sup>23</sup>

In 2005, CHIS administered a Family History of Cancer module to adult participants younger than 65 years ( $n = 33,187$ ).<sup>24</sup> Respondents were asked about any cancer history of first-degree (parents, siblings, children) and second-degree (grandparents, uncles, aunts) blood relatives, including half brothers and sisters. For each female relative diagnosed with cancer, 2 additional questions were asked: 1) whether the female relative had breast, ovarian, uterine, colon, or rectum cancer; and 2) whether the female relative was diagnosed with any of these specific cancers before age 50 years. For each male relative diagnosed with cancer, 2 additional questions were asked: 1) whether the male relative had prostate, colon, rectum, or breast cancer; and 2) whether the male relative was diagnosed with any of these specific cancers before age 50 years.

### **Study Population and Outcome Variables**

The study assessed BC screening adherence and CRC screening adherence separately. The BC screening population was defined as women aged 40 to 64 years with no personal history of BC or ovarian cancer, and who received a mammogram within the past 2 years (total:  $n = 11,885$ ; with a family history:  $n = 1884$ ; without a family history:  $n = 10,001$ ). We included ovarian cancer in the family history of BC group because a family history of ovarian cancer may also increase a woman's risk of BC.<sup>25</sup> Studies on BC screening behavior have included a family history of ovarian cancer as a risk factor for BC risk.<sup>17,26</sup>

The colorectal cancer screening population was defined as adults aged 50 to 64 years with no personal history of colon cancer, who received a fecal occult blood test (FOBT) in the past year, a sigmoidoscopy within the past 5 years, or a colonoscopy within the past 10 years (total:  $n = 11,988$ ; with family history:  $n = 1175$ ; without family history:  $n = 10,813$ ). We could not identify individuals with a personal history of rectal cancer from the CHIS public use data, so only personal history of colon cancer was excluded. For both analyses, the lower age limits were based on the US Preventive Services Task Force guidelines

**Table 1.** CDC Risk Categories Based on Family History

CDC Risk Category	Family History Stratification	BC, <sup>a</sup> Women, Aged 40-64 Years, n = 11,885			CRC, <sup>b</sup> Adults, Aged 50-64 Years, n=11,988		
		No.	% by CDC Risk Category	BC Screening Rates <sup>c</sup> by CDC Risk Category, % (95% CI)	No.	% by CDC Risk Category	CRC Screening Rates <sup>d</sup> by CDC Risk Category, % (95% CI)
Average	No family history	10,001	85.99	76.0 (74.7-77.3)	10,813	91.6	50.5 (49.5-51.6)
Above average	Family history	1884	14.1	83.5 (81.0-85.9)	1175	8.4	71.2 (67.3-75.0)

Abbreviations: BC, breast cancer; CDC, colorectal cancer.

Frequency and rates are weighted estimates.

<sup>a</sup>No personal history of breast or ovarian cancer.

<sup>b</sup>No personal history of colon cancer.

<sup>c</sup>Mammography in past 2 years.

<sup>d</sup>Fecal occult blood testing in past year, sigmoidoscopy in past 5 years, or colonoscopy in past 10 years.

for routine BC and CRC screening at the time the CHIS data was collected in 2005, before the recent changes in the mammogram guideline recommendations in 2009.<sup>27-29</sup>

Although individuals with a family history risk of cancer may be recommended to begin routine cancer screening at earlier ages than the screening guidelines for average-risk individuals, we conducted our analysis on the recommended screening age group for average-risk individuals (aged 40 years and older for mammograms; aged 50 years and older for FOBT, sigmoidoscopy, or colonoscopy) for 2 reasons: 1) to identify subgroups among individuals with a family history who are not even receiving screening per the average-risk guidelines, and 2) to assess racial/ethnic screening disparities among individuals with a family history compared with those with no family history.

### **Risk Stratification by Family History of Cancer**

We categorized family history of either cancers (breast/ovarian or colon) as average risk (no family history) or above average risk (with family history) based on an algorithm developed by Scheuner et al.<sup>30</sup> This risk stratification algorithm is suitable for our data and our analysis, as it employs both first-degree and second-degree relative information and does not require clinical information typically absent in health surveys. Average-risk individuals were those with no family history of cancer (breast/ovarian or colon/rectum) or only 1 second-degree relative diagnosed at any age. Above average risk of family history was defined otherwise. We initially classified the above average-risk category into moderate and high risk, but because of sample size limitations, these 2 risk groups were combined in the study analyses. Screening rates for the moderate and high-risk groups were not statistically different (Table 1).

### **Independent Variables**

Our independent variable of interest was race/ethnicity, specified using the University of California at Los Angeles Center for Health Policy Research classification that codes Latino ethnicity and major race categories as mutually exclusive (white, Latino, Asian/Pacific Islander, African American, and other single race or multiracial). Thus, all the racial/ethnic categories other than Latino are deemed as non-Latino; for example whites are non-Latino whites. In multivariate analyses, we included predictors based on the Andersen Behavioral Model<sup>31</sup> and prior literature assessing screening behavior in multiethnic samples.<sup>7,10,13,32-39</sup> Individual predisposing factors included income status as a percentage of the federal poverty level (FPL) (0%-99%, 100%+), age (continuous in years), sex (for the CRC screening model), marital status (married vs never married, divorced, or widowed), education status (no formal education or less than high school graduate vs high school graduate), limited English proficiency (speaks English less than well vs native speaker and speaks English well or very well), foreign-born versus US-born, and rural versus urban residence. Enabling characteristics were specified as having no health insurance for all or part of the past year and not having paid a visit to the physician in the past 12 months. Need was specified as self-rated health status (fair, poor health vs good, very good, excellent health) and number of chronic conditions (from among the following 7 conditions: diabetes, heart disease, hypertension, asthma, cancer, arthritis, and epilepsy).

### **Statistical Analysis**

We evaluated use of recommended BC and CRC screening separately in 2 study populations: 1) individuals reporting a family history of the cancer associated with the

**Table 2.** Sample Characteristics of Study Populations

Characteristic	Population Recommended for BC Screening, Women, Aged 40-64 Years				Population Recommended for CRC Screening, Adults, Aged 50-64 Years			
	Family History Risk, n = 1884		No Family History Risk, n = 10,001		Family History Risk, n = 1175		No Family History Risk, n = 10,813	
	Weighted % or Mean	No.	Weighted % or Mean	No.	Weighted % or Mean	No.	Weighted % or Mean	No.
Female	100%	1884	100%	10,001	59% <sup>a</sup>	759	51%	6318
Age, mean y (median, SD)	51.2 <sup>a</sup> (51, 7.0)	1884	50.2 (49, 6.9)	10,001	56.6 <sup>a</sup> (56, 4.2)	1,175	56.2 (57, 4.2)	10,813
<b>Race/ethnicity*</b>								
White	70% <sup>a</sup>	1532	55%	6762	75% <sup>a</sup>	978	62%	7993
Latino	12% <sup>a</sup>	116	20%	1290	8% <sup>a</sup>	55	16%	1004
Asian/Pacific Islander	7% <sup>a</sup>	86	14%	1027	7% <sup>a</sup>	59	13%	943
African American	7%	80	7%	513	7%	52	6%	494
Other race	3%	70	4%	409	3%	31	3%	379
Income status as % FPL, 0-99% FPL	6% <sup>a</sup>	100	12%	944	6%	67	9%	789
Education status <high school	7% <sup>a</sup>	84	17%	959	7% <sup>a</sup>	53	13%	862
Single/not married	33% <sup>a</sup>	774	31%	3862	34% <sup>a</sup>	475	30%	4135
Rural	15% <sup>a</sup>	419	12%	1846	15%	261	13%	2183
Limited English proficiency	11% <sup>a</sup>	117	26%	1714	9% <sup>a</sup>	73	22%	1494
Foreign born	18% <sup>a</sup>	234	35%	2412	14% <sup>a</sup>	122	29%	2092
Insured only part/none of past year	12% <sup>a</sup>	222	17%	1488	10% <sup>a</sup>	111	15%	1476
No physician's visit in past year	11%	195	12%	1117	10% <sup>a</sup>	117	14%	1398
Self-rated health: fair/poor	19% <sup>a</sup>	333	23%	1956	21%	229	23%	2174
Chronic conditions, mean of 0-7 (median, SD)	1.02 <sup>a</sup> (1.00, 1.12)	1884	0.91 (1.00, 1.00)	10,001	1.28 <sup>a</sup> (1.00, 1.20)	1175	1.16 (1.00, 1.16)	10,813

Abbreviations: BC, breast cancer; CRC, colorectal cancer; FPL, federal poverty level; SD, standard deviation.

<sup>a</sup>P < .05, proportion of group with family history significantly differs from group with no family history.

\*White, Asian/Pacific Islander, African American and Other race are non-Latino.

screening test, and 2) individuals reporting no family history of the cancer associated with the screening test. We obtained weighted estimates for each study population using sample weights provided in the CHIS public use files. Average-risk (no family history) and above average-risk (no family history) sample distributions were compared using 2-sample test of proportions. By using weighted multivariate logistic regression, we constructed odds ratios and confidence intervals. *F* tests of joint significance with a Bonferroni adjustment were conducted for the race/ethnicity categorical variable. Statistical significance was assessed with a 2-tailed test and alpha of .05. Furthermore, to evaluate whether racial/ethnic disparities significantly varied by family history risk, we estimated models that included all individuals eligible for screening with family history risk, then tested the significance of the interaction effect of race/ethnicity and family history. Data management, variable construction, and regression modeling were conducted using Stata version 10.1 (StataCorp, College Station, Tex).

## RESULTS

A greater proportion of white, higher-income individuals and those with more education were represented in the

population reporting a family history compared with the population reporting no family history for both the BC and CRC screening populations (Table 2). Regardless of risk stratification, whites and individuals with incomes >100% FPL comprised the majority of both the BC and CRC screening populations.

### Breast Cancer Screening

We found no significant racial/ethnic, income, or education disparities in mammography use among women in both (ie, above average risk vs average-risk) family history risk groups, controlling for other characteristics in our models (Table 3). Factors that were significant in predicting BC screening among the group reporting no family history risk were mirrored and amplified in the group reporting a family history risk. These factors included marital status, insurance status, annual physician's visit, self-rated health, and age. Interestingly, rural status, which was not significant in the group with no family history, was significant for the group with a family history (Table 3).

### Colorectal Cancer Screening

Race/ethnicity was significant for both family history risk stratifications (Table 4). However, the racial/ethnic

**Table 3.** Mammography Use in Women, Aged 40 to 64 Years

Characteristic	Family History Risk for Breast or Ovarian Cancer				No Family History for Breast or Ovarian Cancer			
	OR	P	Unweighted, n=1884; Weighted, n = 770,000		OR	P	Unweighted, n=10,001; Weighted, n=4,660,000	
<b>Race/ethnicity (ref: white)</b>			F statistic = 0.14				F statistic = 0.09	
Latino	0.89	.81	0.35	2.25	1.04	.76	0.81	1.33
Asian/Pacific Islander	1.18	.73	0.47	2.98	0.83	.20	0.62	1.11
African American	1.88	.13	0.83	4.24	1.36	.06	0.99	1.85
Other race	0.58	.15	0.28	1.22	0.90	.52	0.65	1.25
Income as % FPL (ref: $\geq 100\%$ FPL), 0-99% FPL	0.55	.13	0.25	1.20	0.81	.09	0.63	1.03
Education (ref: high school graduate), <high school	0.98	.98	0.35	2.80	0.92	.58	0.68	1.24
Single/not married	0.59 <sup>a</sup>	.03	0.37	0.95	0.71 <sup>a</sup>	<.01	0.61	0.83
Rural	0.58 <sup>a</sup>	.03	0.35	0.95	0.90	.28	0.73	1.10
Limited English proficiency	1.96	.26	0.61	6.31	1.06	.75	0.76	1.47
Foreign born	2.00	.13	0.82	4.85	1.11	.43	0.86	1.43
Insured part/none of past year	0.25 <sup>a</sup>	<.01	0.15	0.43	0.45 <sup>a</sup>	<.01	0.37	0.56
No physician's visit in past year	0.19 <sup>a</sup>	<.01	0.12	0.29	0.32 <sup>a</sup>	<.01	0.27	0.37
Self-rated health: fair/poor	0.55 <sup>a</sup>	.02	0.33	0.92	0.77 <sup>a</sup>	.05	0.59	1.00
Chronic conditions	1.04	.66	0.88	1.22	1.07	.14	0.98	1.18
Age	1.08 <sup>a</sup>	<.01	1.05	1.11	1.08 <sup>a</sup>	<.01	1.07	1.10

Abbreviations: CI, confidence interval; FPL, federal poverty level; OR, odds ratio.

<sup>a</sup>Significant at  $P < .05$ ,  $F$  test of joint significance of race/ethnicity with Bonferroni adjustment.

disparities (Latinos, Asians and Pacific Islanders, other race compared with whites) seen in the group with no family history risk were only evident among Latinos compared with whites. Latinos reporting no family history had  $0.74\times$  the odds of being screened for CRC compared with whites. This gap was more substantial among the group with family history risk (odds ratio [OR], 0.28; 95% confidence interval [CI], 0.11-0.60). On the basis of a model testing the interaction term of Latino and family history (data not shown), the difference in the magnitude of the Latino-white disparity between those with and without a family history was statistically significant ( $P < .05$ ). The sex disparity seen in the group with no family history was not significant in the group reporting a family history of CRC. Lacking insurance and not having an annual physician visit were significantly associated with CRC screening in both family history risk categories.

## DISCUSSION

Our study had 3 major findings. First, in California, where mammography rates are high,<sup>40</sup> we found no significant disparities by race/ethnicity and income in both family history risk groups. Although our study may not be generalizable nationally, this finding is consistent with Wu et al's study using the National Health Interview Survey.<sup>41</sup> Second, racial/ethnic disparities were more evident in CRC screening, and we found evidence of a Latino-

white gap among the group with a family history. As this is the new contribution of the study, these results should be further explored and confirmed in national population-based data and in medical claims data that contain information on family history. Third and most importantly, personal knowledge of family history did not close the Latino-white gap in CRC screening. This finding is relevant and new, not found in previous diverse population-based studies that have included Latinos with a family history.<sup>5,41</sup> Our estimate that Latinos with a family history risk had  $0.28\times$  the odds of being screened for CRC compared with whites points to a considerable disparity, greater than most of the detected disparities from other factors in our model, and of greater magnitude than the Latino-white difference among individuals with no family history (OR, 0.74; 95% CI, 0.59-0.92).

There are several explanations regarding why Latinos are not getting screened for CRC as much as whites, even if they know that they have a family member with CRC. On the patient side, the most relevant study to put our results in context is the recent national population-based study conducted by Orom et al using the national 2007 Health Information Trends Survey. In that study of a multiethnic sample, Orom and her colleagues found 1) that Latinos had lower perceived cancer risk than whites, 2) that the lower rates of perceived cancer risk were associated with lower rates of reported family history of cancer among Latinos compared with whites, and most

**Table 4.** Colorectal Cancer Screening, Adults Aged 50 to 64 Years

Characteristic	Family History Risk for Colorectal Cancer				No Family History for Colorectal Cancer			
	Unweighted, n=1175; Weighted, n=460,000				Unweighted, n=10,813; Weighted, n=4,990,000			
	OR	P	95% CI		OR	P	95% CI	
Female	0.85	.44	0.59	1.32	0.83 <sup>a</sup>	<.01	0.75	0.93
<b>Race/ethnicity (ref: white)</b>			<i>F</i> statistic = 0.05				<i>F</i> statistic < 0.01	
Latino	0.28 <sup>a</sup>	.01	0.11	0.60	0.74 <sup>a</sup>	.01	0.59	0.92
Asian/Pacific Islander	0.68	.35	0.40	2.12	0.74 <sup>a</sup>	.01	0.59	0.94
African American	0.92	.84	0.31	1.34	1.08	.53	0.84	1.40
Other race	0.47	.21	0.16	1.84	0.66 <sup>a</sup>	.03	0.46	0.95
Income as % FPL (ref: ≥100% FPL), 0-99% FPL	0.49	.06	0.17	0.90	0.83	.15	0.65	1.07
Education (ref: high school graduate), <high school	1.01	.99	0.34	3.25	0.80	.14	0.59	1.08
Single/not married	0.82	.31	0.56	1.21	0.81 <sup>a</sup>	<.01	0.73	0.90
Rural	1.36	.21	0.84	2.19	0.89	.20	0.75	1.06
Limited English proficiency	1.14	.85	0.30	4.38	0.79	.19	0.55	1.12
Foreign born	1.35	.44	0.62	2.94	1.03	.79	0.85	1.24
Insured part/none of past year	0.25 <sup>a</sup>	<.01	0.14	0.45	0.45 <sup>a</sup>	<.01	0.37	0.54
No physician's visit in past year	0.33 <sup>a</sup>	<.01	0.17	0.63	0.29 <sup>a</sup>	<.01	0.25	0.35
Self-rated health: fair/poor	0.92	.73	0.58	1.46	1.02	.86	0.86	1.20
Chronic conditions	1.05	.60	0.87	1.26	1.08 <sup>a</sup>	.01	1.02	1.15
Age	1.13 <sup>a</sup>	<.01	1.07	1.19	1.10 <sup>a</sup>	<.01	1.08	1.11

Abbreviations: CI, confidence interval; FPL, federal poverty level; OR, odds ratio.

<sup>a</sup>Significant at  $P < .05$ , *F* test of joint significance of race/ethnicity with Bonferroni adjustment.

importantly, 3) that reporting a family history of cancer was positively associated with higher perceived cancer risk among whites, but not among Latinos. Their last finding supports our study results of the widening Latino-white disparity in CRC screening among the group with a family history of CRC. The authors further posit that Latinos with a family history may be less aware of their cancer risk “due to language and other barriers that can make the dissemination of health information difficult.”<sup>14</sup>

Even among Latinos who do perceive their family history of colorectal cancer to be associated with increased risk of colorectal cancer, some studies have suggested that compared with whites, on average, Latinos may harbor greater fear or denial of this risk, causing them to delay or fail to seek colorectal cancer screening.<sup>16,17,32</sup> However, in a study with a diverse sample of women recruited from primary care clinics in San Francisco (42% reported having a family history of cancer), among Latinas who had the highest perceived risk of 3 cancers, including colon cancer, their perceived risk was associated with obtaining cancer screening tests.<sup>42</sup> The study's participants, women associated with a primary care clinic, may be a select sample of women who tend to value preventive care and who have a regular provider. Thus, they may be more motivated to be screened for CRC than the average Latino with a family history of CRC. Nevertheless, interventions

are needed that address the spectrum of reactions to knowledge of a family history of CRC (ie, fear, denial, anxiety, and indifference) among affected Latinos to mobilize CRC screening behavior.

On the provider side, studies suggest that rates of family history inquiries are low in routine clinical encounters.<sup>13,43</sup> We found no studies that determined whether there is a differential rate of obtaining family histories by race/ethnicity. We therefore posit that because obtaining family history and explaining increased risks and recommended screening intervals may require more time in the patient-physician encounter, language and cultural barriers may, on average, differentially deter this knowledge transfer of CRC risk to Latinos but not to whites. In Guerra et al's qualitative study on barriers and facilitators to physician recommendation of CRC screening, physicians caring for non-English-speaking patients reported that they “had a particularly difficult time recommending CRC screening because translation of the recommendation takes up much of the time allotted for the visit.”<sup>44</sup> Similarly, Wee et al's study found that Latino adults are less likely to receive counseling from their physicians about CRC tests than whites.<sup>21</sup> Wee et al's study implies that the Latino-white disparity could be more pronounced among those with a family history, because family history counseling requires an even greater time investment.

The differential knowledge transfer of CRC risk may also result when the patient-provider interaction is racially/ethnically discordant. For example, Ge et al 2009 found that “physicians did not solicit or address cultural barriers to CRC screening and patients did not volunteer culture-related concerns regarding CRC screening” in ethnically discordant physician-patient interactions.<sup>45</sup> Drawing on Guerra et al’s study previously discussed,<sup>44</sup> because counseling for family history risk is more complex than average-risk screening recommendations and would require more time, the Latino disadvantages in linguistically and culturally discordant physician relationships may be greater in individuals with a family history of CRC. Discordant ethnic patient-physician relationships are certainly prevalent in California, as Latinos are more than a third of California’s population, but make up only 5% of the state’s physician population.<sup>46</sup> Thus, racial/ethnic variations in the provider-patient interaction could be a source of the CRC screening disparity, and has a potentially greater penalizing effect on Latinos with a family history of CRC. Finally, our results in California are likely to be present for Latinos across the United States, and could be more pronounced in areas with fewer cultural and linguistic services for Latinos.

A key limitation of our study is that it is based on self-reported survey data. Family history reports appear to be generally accurate; validation studies on self-reports of cancer family history suggest reports of first-degree relatives to be highly accurate<sup>2,47-49</sup> and reports of second-degree relatives to be moderately accurate. One study suggests that the validity of self-reported family history is better for BC and CRC than ovarian and endometrial cancers. Although self-reported family history is generally valid, recent meta-analysis on the accuracy of self-reported cancer screening suggests screening use in population-based surveys tends to be over-reported,<sup>50</sup> especially among ethnic minority respondents; thus, the racial/ethnic disparities we report here may be underestimated. The sample size of minority racial/ethnic groups in the CHIS family history population may also have limited our capability to detect any disparities in screening by population subgroups. However, we detected a significant effect for Latinos, whose sample size was comparable to that of Asians and Pacific Islanders and African Americans. Finally, our study approach evaluated screening beginning at the age recommended for average-risk adults to compare across all risk categories. Because of smaller samples, we could not ascertain whether disparities may be narrower or wider in evaluating adults with a family

history who received mammograms at age <40 years and CRC screening at age <50 years.

Despite these limitations, our study provides a compelling picture of how a family history analysis could direct where efforts are most needed in reducing cancer disparities by race and ethnicity.<sup>51,52</sup> Among California adults, our troubling finding was that knowledge of their family history of CRC did not close the Latino-white gap in CRC screening, but actually widened the disparity. More aggressive interventions that enhance the communication between Latinos and their physicians about family history and cancer risk are needed and could reduce the substantial Latino-white screening disparity in Latinos most susceptible to CRC.<sup>51,52</sup>

## FUNDING SOURCES

This study was supported by the University of California at San Francisco Center for Translational and Policy Research on Personalized Medicine (TRANSPERS)-National Cancer Institute (NCI) P01 CA 130818 (principal investigator [PI], K. Phillips), NCI K07 CA100097 (PI, N.A.P.), Network for Multicultural Research on Health and Healthcare, Department of Family Medicine, David Geffen School of Medicine, University of California at Los Angeles (UCLA), funded by the Robert Wood Johnson Foundation, and UCLA Summer Graduate Research Mentorship Program. This research was also conducted with support from the Health Disparities Research Program of Harvard Catalyst/Harvard Clinical and Translational Science Center (NIH award #UL1 RR 025758 to J.S.H. and financial contributions from Harvard University and its affiliated academic health care centers).

## CONFLICT OF INTEREST DISCLOSURES

The authors made no disclosures.

## REFERENCES

1. Noe M, Schroy P, Demierre MF, Babayan R, Geller AC. Increased cancer risk for individuals with a family history of prostate cancer, colorectal cancer, and melanoma and their associated screening recommendations and practices. *Cancer Causes Control*. 2008;19:1-12.
2. Anton-Culver H, Kurosaki T, Taylor TH, Gildea M, Brunner D, Bringman D. Validation of family history of breast cancer and identification of the BRCA1 and other syndromes using a population-based cancer registry. *Genet Epidemiol*. 1996;13:193-205.
3. Phipps AI, Buist DS, Malone KE, et al. Family history of breast cancer in first-degree relatives and triple-negative breast cancer risk. *Breast Cancer Res Treat*. 2011;126:671-678.
4. Rim SH, Seff L, Ahmed F, King JB, Coughlin SS. Colorectal cancer incidence in the United States, 1999-2004: an updated analysis of data from the National Program of Cancer Registries and the Surveillance, Epidemiology, and End Results Program. *Cancer*. 2009;115:1967-1976.
5. Ramsey SD, Yoon P, Moonesinghe R, Khoury MJ. Population-based study of the prevalence of family history of cancer: implications for cancer screening and prevention. *Genet Med*. 2006;8:571-575.

6. Smith-Bindman R, Miglioretti DL, Lurie N, et al. Does utilization of screening mammography explain racial and ethnic differences in breast cancer? *Ann Intern Med.* 2006;144:541-553.
7. Edwards QT, Li AX, Pike MC, et al. Ethnic differences in the use of regular mammography: the multiethnic cohort. *Breast Cancer Res Treat.* 2009;115:163-170.
8. James TM, Greiner KA, Ellerbeck EF, Feng C, Ahluwalia JS. Disparities in colorectal cancer screening: a guideline-based analysis of adherence. *Ethn Dis.* 2006;16:228-233.
9. Wells KJ, Roetzheim RG. Health disparities in receipt of screening mammography in Latinas: a critical review of recent literature. *Cancer Control.* 2007;14:369-379.
10. Burgess DJ, van Ryn M, Grill J, et al. Presence and correlates of racial disparities in adherence to colorectal cancer screening guidelines. *J Gen Intern Med.* 2011;26:251-258.
11. Wojcik BE, Spinks MK, Stein CR. Effects of screening mammography on the comparative survival rates of African American, white, and Hispanic beneficiaries of a comprehensive health care system. *Breast J.* 2003;9:175-183.
12. Du XL, Lin CC, Johnson NJ, Altekruze S. Effects of individual-level socioeconomic factors on racial disparities in cancer treatment and survival: findings from the National Longitudinal Mortality Study, 1979-2003. *Cancer.* 2011;117:3242-3251.
13. Karliner LS, Napoles-Springer A, Kerlikowske K, Haas JS, Gregorich SE, Kaplan CP. Missed opportunities: family history and behavioral risk factors in breast cancer risk assessment among a multiethnic group of women. *J Gen Intern Med.* 2007;22:308-314.
14. Orom H, Kiviniemi MT, Underwood W III, Ross L, Shavers VL. Perceived cancer risk: why is it lower among nonwhites than whites? *Cancer Epidemiol Biomarkers Prev.* 2010;19:746-754.
15. Stacy R, Torrence WA, Mitchell CR. Perceptions of knowledge, beliefs, and barriers to colorectal cancer screening. *J Cancer Educ.* 2008;23:238-240.
16. Walsh JM, Kaplan CP, Nguyen B, Gildengorin G, McPhee SJ, Perez-Stable EJ. Barriers to colorectal cancer screening in Latino and Vietnamese Americans compared with non-Latino white Americans. *J Gen Intern Med.* 2004;19:156-166.
17. Andersen MR, Smith R, Meischke H, Bowen D, Urban N. Breast cancer worry and mammography use by women with and without a family history in a population-based sample. *Cancer Epidemiol Biomarkers Prev.* 2003;12:314-320.
18. Kagawa-Singer M, Dadia AV, Yu MC, Surbone A. Cancer, culture, and health disparities: time to chart a new course? *CA Cancer J Clin.* 2010;60:12-39.
19. Pasick RJ, Barker JC, Otero-Sabogal R, Burke NJ, Joseph G, Guerra C. Intention, subjective norms, and cancer screening in the context of relational culture. *Health Educ Behav.* 2009;36(5 suppl):91S-110S.
20. Rich SE, Kuyateh FM, Dwyer DM, Groves C, Steinberger EK. Trends in self-reported health care provider recommendations for colorectal cancer screening by race. *Prev Med.* 2011;53:70-75.
21. Wee CC, McCarthy EP, Phillips RS. Factors associated with colon cancer screening: the role of patient factors and physician counseling. *Prev Med.* 2005;41:23-29.
22. California Health Interview Survey: CHIS 2005 Methodology Series: Report 1-Sample Design. Los Angeles, CA: UCLA Center for Health Policy Research; 2007.
23. Prograis LJ Jr, Zunich KM. NIAID programs for asthma research, education, and outreach. *Chest.* 1992;101(6 suppl):357S-358S.
24. California Health Interview Survey: CHIS 2005 Family Cancer History Supplement. Los Angeles, CA: UCLA Center for Health Policy Research; 2007.
25. Antoniou A, Pharoah PD, Narod S, et al. Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected in case series unselected for family history: a combined analysis of 22 studies. *Am J Hum Genet.* 2003;72:1117-1130.
26. Halbert CH, Kessler L, Wileyto EP, et al. Breast cancer screening behaviors among African American women with a strong family history of breast cancer. *Prev Med.* 2006;43:385-388.
27. Screening for colorectal cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med.* 2008;149:627-637.
28. Screening for breast cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med.* 2009;151:716-726, W-236.
29. Whitlock EP, Lin JS, Liles E, Beil TL, Fu R. Screening for colorectal cancer: a targeted, updated systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med.* 2008;149:638-658.
30. Scheuner MT, Wang SJ, Raffel LJ, Larabell SK, Rotter JI. Family history: a comprehensive genetic risk assessment method for the chronic conditions of adulthood. *Am J Med Genet.* 1997;71:315-324.
31. Andersen RM. Revisiting the behavioral model and access to medical care: does it matter? *J Health Soc Behav.* 1995;36:1-10.
32. Haas JS, Kaplan CP, Des Jarlais G, Gildengoin V, Perez-Stable EJ, Kerlikowske K. Perceived risk of breast cancer among women at average and increased risk. *J Womens Health (Larchmt).* 2005;14:845-851.
33. Slomiany BA, McMasters KM, Chagpar AB. The recent decline in mammography rates is limited to low- to average-risk women. *Am J Surg.* 2008;196:821-826; discussion 26.
34. Laing SS, Makambi K. Predicting regular breast cancer screening in African-American women with a family history of breast cancer. *J Natl Med Assoc.* 2008;100:1309-1317.
35. Williams KP, Sheppard VB, Todem D, Mabiso A, Wulu JT Jr, Hines RD. Family matters in mammography screening among African-American women age > 40. *J Natl Med Assoc.* 2008;100:508-515.
36. Griffith KA, McGuire DB, Royak-Schaler R, Plowden KO, Steinberger EK. Influence of family history and preventive health behaviors on colorectal cancer screening in African Americans. *Cancer.* 2008;113:276-285.
37. Ponce NA, Huh S, Bastani R. Do HMO market level factors lead to racial/ethnic disparities in colorectal cancer screening? A comparison between high-risk Asian and Pacific Islander Americans and high-risk whites. *Med Care.* 2005;43:1101-1108.
38. Kagawa-Singer M, Pourat N, Breen N, et al. Breast and cervical cancer screening rates of subgroups of Asian American women in California. *Med Care Res Rev.* 2007;64:706-730.
39. Etzioni DA, Ponce NA, Babey SH, et al. A population-based study of colorectal cancer test use: results from the 2001 California Health Interview Survey. *Cancer.* 2004;101:2523-2532.
40. Kaiser Family Foundation. California: Percent of women age 40 and older who report having had a mammogram within the last 2 years, 2010. Available at: <http://www.statehealthfacts.org/comparemaptable.jsp?cat=10&ind=479>. Accessed on August 8, 2011.
41. Wu H, Zhu K, Jatoti I, Shah M, Shriver CD, Potter J. Factors associated with the incompletion with mammogram screening

- among individuals with a family history of breast cancer or ovarian cancer. *Breast Cancer Res Treat.* 2007;101:317-324.
42. Kim SE, Perez-Stable EJ, Wong S, et al. Association between cancer risk perception and screening behavior among diverse women. *Arch Intern Med.* 2008;168:728-734.
  43. Fletcher RH, Lobb R, Bauer MR, et al. Screening patients with a family history of colorectal cancer. *J Gen Intern Med.* 2007;22:508-513.
  44. Guerra CE, Schwartz JS, Armstrong K, Brown JS, Halbert CH, Shea JA. Barriers of and facilitators to physician recommendation of colorectal cancer screening. *J Gen Intern Med.* 2007;22:1681-1688.
  45. Ge G, Burke N, Somkin CP, Pasick R. Considering culture in physician-patient communication during colorectal cancer screening. *Qual Health Res.* 2009;19:778-789.
  46. Bates T, Chapman S. Diversity in California's health professions: physicians. A report of the California Health Workforce Tracking Collaborative. UCSF Center for the Health Professions, San Francisco, CA: University of California at San Francisco; 2008.
  47. Parent ME, Ghadirian P, Lacroix A, Perret C. Accuracy of reports of familial breast cancer in a case-control series. *Epidemiology.* 1995;6:184-186.
  48. Ziogas A, Anton-Culver H. Validation of family history data in cancer family registries. *Am J Prev Med.* 2003;24:190-198.
  49. Murff HJ, Byrne D, Syngal S. Cancer risk assessment: quality and impact of the family history interview. *Am J Prev Med.* 2004;27:239-245.
  50. Rauscher GH, Johnson TP, Cho YI, Walk JA. Accuracy of self-reported cancer-screening histories: a meta-analysis. *Cancer Epidemiol Biomarkers Prev.* 2008;17:748-757.
  51. Qureshi N, Wilson B, Santaguada P, et al. Family history and improving health. *Evid Rep Technol Assess (Full Rep).* 2009;(186):1-135.
  52. Berg AO, Baird MA, Botkin JR, et al. National Institutes of Health State-of-the-Science Conference Statement: Family History and Improving Health: August 24-26, 2009. *NIH Consens State Sci Statements.* 2009;26:1-19.